

EHC ED Critical Care Induced Hypothermia Protocol

Name: _____

Place Sticker

MRN: _____

Date: _____ Time of Screening: _____

Inclusion Criteria (Must have All)

- ☐ Post Cardiac Arrest (Any rhythm as cause of arrest is eligible)
- ☐ ROSC < 30 min from EMS/Code Team Arrival
- ☐ Time now < 6 hrs from ROSC
- ☐ Comatose (Does not follow commands)
- ☐ MAP > 65 on no more than one vasopressor

Exclusion Criteria

- ☐ Pt has DNR, MOLST, **poor baseline status**, or terminal disease
- ☐ Active or Intracranial Bleeding
- ☐ Traumatic etiology for arrest
- ☐ Cryoglobulinemia
- ☐ Pregnancy (Relative-Consider OB/Gyn consult)
- ☐ Recent Major Surgery (Relative)
- ☐ Sepsis as cause of Arrest (Relative)

Neurologic Exam

Eye Opening

Spontaneous ----- ☐ 4
Voice ----- ☐ 3
Pain ----- ☐ 2
None ----- ☐ 1

Verbal

Oriented ----- * ☐ 5
Confused ----- * ☐ 4
Inappropriate ----- ☐ 3
Sounds ----- ☐ 2
None ----- ☐ 1
Intubated ----- ☐ 1

Motor

Obeys ----- * ☐ 6
Localizes ----- ☐ 5
Withdraws ----- ☐ 4
Decorticate ----- ☐ 3
Decerebrate ----- ☐ 2
None ----- ☐ 1

Brainstem

Pupils React ☐ yes ☐ no
Corneal ☐ yes ☐ no
Spontaneous
Respirations ☐ yes ☐ no
Doll's Eyes ☐ yes ☐ no

DTRs:

Bicep L R

Knee L R

Toes L R

List any Sedatives or Paralytics On-Board at time of Exam: _____

If any Starred (*) Item is checked off on the neuro exam, the patient is ineligible for the protocol.

Protocol

- Discuss Case with ICU Fellow or Attending (They must agree with the plan for hypothermia)
- Time of Discussion: _____ If pt is deemed ineligible by ICU, list reason: _____
- List Initial Arrest Rhythm: _____ List Number of Minutes from Start of CPR to ROSC: _____
- Send blood for: CMP, LFTs, Superstat I, Lactate, CBC, PT/PTT, CK/MB/Troponin, Lipase/Amylase
- Place foley catheter and monitor urine output.
- Completely expose patient and place cooling blanket above and below with nothing between blanket & skin.
- Place temp probe in **mid-esophagus** (~4 cm above xiphoid via oral/nasal); if unable to place in esophagus, probe can be placed rectally (5 cm)
- Hook both cooling blankets and the probe to the same blanketrol machine.
- Set temperature to 33° C and Set the machine to "**Auto Control**".
- List time Now: _____ List Initial Patient Temperature: _____° C
- If initial temperature is < 33° C, allow patient to warm to 33° C.
- Begin opioids & sedation. Titrate to Ramsay Score 4/5.
- Infuse refrigerated crystalloid, preferably through large bore, peripheral IV.
Administer at ~100 ml per minute using pressure bag (evacuate air first). Maximum initial infusion is 30 cc/kg; if patient not < 34° C after this amount, wait 15 minutes before giving further 250 cc boluses Q 10 minutes.
- Administer Tylenol 650 mg GT Q 6 hours unless pt has allergy.
- If during induction, pt has shivering unrelieved by the above meds, Vecuronium 0.1 mg/kg x1 can be used
- Total Cold Crystalloid Infused: _____ Time that Pt reaches 34° C: _____
- If patient's temperature rises above 34.5° C, infuse 250 cc boluses of cold crystalloid Q 10 min until < 34° C.
- Assess for shivering Q 15 minutes. If any signs of shivering, see the protocol on page 4.
- Maintain temperature 32-34° C for 24 hours (ideal temperature is 33° C).
- If significant bleeding or severe hemodynamic instability, consider rewarming. **See ehced.org for protocol.**
- Time of Rewarming: _____ Reason Necessary: _____
- Maintain MAP > 80: Pressors and/or Dobutamine may be used during protocol, if fluid loading ineffective.

Post-ROSC Care Package

Induction of Hypothermia

See First Page

Procedures

- Full sterile neck line with CVP monitoring
- Full sterile femoral arterial line (Axillary if femoral contraindicated/unsuccessful)
- Foley Catheter with hourly urine monitoring
- Orogastric Tube on suction

Ventilation

- Place patient on AC Mode
- Set Vt to 8 ml/kg IBW (see last page)
- Set IFR to 60 lpm
- Set Initial rate to 18 bpm
- Set Initial O2 to 50%
- Titrate FiO2/PEEP to achieve corrected ABG Saturation 94-96%.
- Often pulse ox will not read well due to peripheral vasoconstriction
- Send an ABG, **DO NOT INDICATE THE PATIENT'S TEMPERATURE ON THE ABG ORDER**

Hemodynamic Goals

- **Ensure Adequate Preload**
Assess by passive leg raise, pulse pressure variation, and echo. CVP may provide some indication if very low. Use normal saline or lactated ringers boluses. Use room temperature fluid if patient at goal temperature. Replace patient's urine losses 1:1
- **MAP > 65** at all times, MAP > 80 is preferred to augment cerebral perfusion
Preferred initial pressor is norepinephrine, may add vasopressin if necessary
If MAP is < 80 and CVP > 10 perform passive straight leg raise to assess fluid responsiveness.
If MAP > 100, start nitroglycerin infusion
- **Corrected ScvO2 > 70**
Can be measured by PreSEP catheter or central venous O2 saturation (send blood gas as mixed venous)
If ScvO2 < 70 and HB < 7 (some would advocate <10 as trigger), transfuse patient
If HB > 7, evaluate echocardiogram and consider inotropes vs. balloon pump/revascularization
- **Lactate**
Hypothermia will raise lactate levels and post-arrest patients will have high lactate. Send a baseline level after the patient achieves goal temperature. From this point on, the lactate should stay the same or drop. If lactate is increasing, the patient is under-resuscitated or seizing

Sedation

- To gain the full benefits of hypothermia, it is imperative that the patient is adequately sedated
- Optimize fentanyl infusion rate first
- Add on propofol or dexmedetomidine if necessary
- Titrate to Ramsay Score of 4/5 (see last page)

Post-ROSC Care Package

Labs & Electrolytes

- Send Superstat I (ABG with Electrolytes) and Lactate Q 1 hour for first 4 hours, then Q 4 hours
- On arrival, send CMP, CBC, Lytes, PT/PTT, Lipase, Cardiac Enzymes, Type and Hold, & Pan-Cultures
- Send CMP (complete metabolic panel) and CBC Q 4 hours
- Send Cardiac Enzymes Q 6 hours
- Keep Magnesium at high-normal at all times with aggressive IV repletion
- Replete Potassium if < 3.4 with IV KCl
- Keep iCal at high normal at all times
- Keep Sodium at least 140 at all times, 150 is preferable
- Keep Glucose < 150 with Insulin Drip (preferred) or Subcutaneous Regular Insulin

Cardiac Testing

- Get EKG immediately upon arrival; at the start of hypothermia induction; and Q 1 hour for the first 4 hours
- If possible, get a bedside transthoracic echo at the start of induction. In the ED, this should be performed by the emergency physician or cardiology. Look specifically for qualitative LV function, RV function, pericardial effusion/tamponade, & gross valve function

DVT Prophylaxis

- If no contraindication, Heparin 5000 units subcutaneous Q 8 hours

Stress Ulcer Prophylaxis

- Nexium 40 mg IVSS x 1

VAP Prophylaxis

- Head of bed to 30°
- Place in-line closed suction and perform aggressive pulmonary toilet

Additional Testing

- Consider Head CT if possible neurologic cause to arrest. Note: even an intracranial bleed is not a contra-indication to continuation of induced hypothermia. Consider letting the patient drift to 34°C and administration of dDAVP.
- Consider CTA if strong suspicion of PE as the cause of arrest. Bedside dopplers by EP or sono technician may be a good first step
- EEG if seizures (convulsive or non-convulsive) are suspected

Revascularization for STEMI

- PCI is preferred, consult with CPORT fellow/attending and CCU fellow. Hypothermia does not need to be discontinued for PCI
- If PCI is not available or will be delayed, thrombolysis should be administered. Thrombolysis can be given during hypothermia. CPR performed prior to ROSC should not stop reperfusion therapy. Use standard doses of Retevase. Consult with CPORT fellow/attending.

Post-ROSC Care Package

Ramsay Sedation Scale

- 1 Patient is anxious and agitated or restless, or both
- 2 Patient is co-operative, oriented, and tranquil
- 3 Patient responds to commands only
- 4 Patient exhibits brisk response to light glabellar tap or loud auditory stimulus
- 5 Patient exhibits a sluggish response to light glabellar tap or loud auditory stimulus
- 6 Patient exhibits no response

Induced Hypothermia Shivering Protocol

Shivering Protocol After Induction

Bedside Shivering Assessment (BSAS) (Neurocrit Care 2007;6:213)

0-None, no shivering. Must not have shivering on EKG or palpation.

1-Mild-localized to neck/thorax. May only be noticed on palpation or EKG.

2-Moderate-intermittent involvement of upper extremities +/- thorax

3-Severe-generalized shivering or sustained upper extremity shivering

•All patients receive:

Acetaminophen 650 mg GT Q 6 hours unless allergic

•If BSAS > 1, add **Fentanyl Drip** (titrate as per EHCED drip sheet)

•If BSAS still > 1, add **Propofol Drip** (titrate as per EHCED drip sheet)

•If BSAS still > 1, administer **MgSO₄** 2 grams IVSS, then 0.5-1 gram/hr for target serum Mg 3 mg/dl

•If BSAS still > 1, add **Bair Hugger Device** on both of patient's arms

•If BSAS still > 1, administer **Ketamine** 0.5 mg/kg IVP

•If BSAS still > 1 after titration of above meds, add **Nimbex** 0.15 mg/kg IV Q 1 hour PRN

Paralysis should only be necessary under extraordinary circumstances!

ARDSNet Vent Protocol



NIH NHLBI ARDS Clinical Network
Mechanical Ventilation Protocol Summary
www.ardsnet.org

INCLUSION CRITERIA: Acute onset of

1. $\text{PaO}_2/\text{FiO}_2 \leq 300$ (corrected for altitude)
2. Bilateral (patchy, diffuse, or homogeneous) infiltrates consistent with pulmonary edema
3. No clinical evidence of left atrial hypertension

PART I: VENTILATOR SETUP AND ADJUSTMENT

1. Calculate predicted body weight (PBW)
Males = $50 + 2.3 [\text{height (inches)} - 60]$
Females = $45.5 + 2.3 [\text{height (inches)} - 60]$
2. Select Assist Control Mode
3. Set initial TV to 8 ml/kg PBW
4. Reduce TV by 1 ml/kg at intervals ≤ 2 hours until TV = 6 ml/kg PBW.
5. Set initial rate to approximate baseline VE (not > 35 bpm).
6. Adjust TV and RR to achieve pH and plateau pressure goals below.
7. Set inspiratory flow rate above patient demand (usually > 80L/min)

OXYGENATION GOAL: PaO_2 55-80 mmHg or SpO_2 88-95%

Use incremental FiO_2 /PEEP combinations below to achieve goal

FiO₂	0.3	0.4	0.4	0.5	0.5	0.6	0.7	0.7
PEEP	5	5	8	8	10	10	10	12

FiO₂	0.7	0.8	0.9	0.9	0.9	1.0	1.0	1.0
PEEP	14	14	14	16	18	20	22	24

PLATEAU PRESSURE GOAL: 30 cm H₂O

Check Pplat (0.5 second inspiratory pause), SpO_2 , Total RR, TV and pH (if available) at least q 4h and after each change in PEEP or TV.

If Pplat > 30 cm H₂O: decrease TV by 1 ml/kg steps (minimum = 4 ml/kg).

If Pplat < 25 cm H₂O: TV < 6 ml/kg, increase TV by 1 ml/kg until Pplat > 25 cm H₂O or TV = 6 ml/kg.

If Pplat < 30 and breath stacking occurs: may increase TV in 1 ml/kg increments (maximum = 8 ml/kg).

pH GOAL: 7.30-7.45

Acidosis Management: (pH < 7.30)

If pH 7.15-7.30: Increase RR until pH > 7.30 or $\text{PaCO}_2 < 25$ (Maximum RR = 35).

If RR = 35 and $\text{PaCO}_2 < 25$, may give NaHCO_3 .

If pH < 7.15: Increase RR to 35.

If pH remains < 7.15 and NaHCO_3 considered or infused, TV may be increased in 1 ml/kg steps until pH > 7.15 (Pplat target may be exceeded).

Alkalosis Management: (pH > 7.45) Decrease vent rate if possible.